

p -ADIC MULTIDIMENSIONAL WAVELETS AND THEIR APPLICATION TO p -ADIC PSEUDO-DIFFERENTIAL OPERATORS

A. YU. KHRENNIKOV AND V. M. SHELKOVICH

ABSTRACT. In this paper we study some problems related with the theory of multidimensional p -adic wavelets in connection with the theory of multidimensional p -adic pseudo-differential operators (in the p -adic Lizorkin space). We introduce a new class of n -dimensional p -adic compactly supported wavelets. In one-dimensional case this class includes the Kozyrev p -adic wavelets. These wavelets (and their Fourier transforms) form an orthonormal complete basis in $\mathcal{L}^2(\mathbb{Q}_p^n)$. A criterion for a multidimensional p -adic wavelet to be an eigenfunction for a pseudo-differential operator is derived. We prove that these wavelets are eigenfunctions of the Taibleson fractional operator. Since many p -adic models use pseudo-differential operators (fractional operator), these results can be intensively used in applications. Moreover, p -adic wavelets are used to construct solutions of linear and *semi-linear* pseudo-differential equations.

1. INTRODUCTION

There are a lot of papers where different applications of p -adic analysis to physical problems, stochastics, cognitive sciences and psychology are studied [6]–[10], [13]–[19], [30]–[32] (see also the references therein).

The field \mathbb{Q}_p of p -adic numbers is defined as the completion of the field of rational numbers \mathbb{Q} with respect to the non-Archimedean p -adic norm $|\cdot|_p$. This norm is defined as follows: $|0|_p = 0$; if an arbitrary rational number $x \neq 0$ is represented as $x = p^\gamma \frac{m}{n}$, where $\gamma = \gamma(x) \in \mathbb{Z}$, and m and n are not divisible by p , then $|x|_p = p^{-\gamma}$. This norm in \mathbb{Q}_p satisfies the strong triangle inequality $|x + y|_p \leq \max(|x|_p, |y|_p)$.

It is known that for the p -adic analysis related to the mapping $\mathbb{Q}_p \rightarrow \mathbb{C}$, where \mathbb{C} is the field of complex numbers, the operation of partial differentiation is *not defined*, and as a result, large number of models connected with p -adic

Date:

2000 *Mathematics Subject Classification.* Primary 11F85, 42C40, 47G30; Secondary 26A33, 46F10.

Key words and phrases. p -adic compactly supported wavelets; p -adic pseudo-differential operators; fractional operators; p -adic Lizorkin spaces.

This paper was supported in part by the grant of The Swedish Royal Academy of Sciences on collaboration with scientists of former Soviet Union. The second author (V. S.) was also supported in part by DFG Project 436 RUS 113/809 and Grant 05-01-04002-NNIOa of Russian Foundation for Basic Research.

differential equations use pseudo-differential operators and the theory of p -adic distributions (generalized functions) (see the above mentioned papers and books). In particular, fractional operators $D^\alpha = f_{-\alpha}*$ are extensively used, where f_α is the p -adic *Riesz kernel*, $*$ is a convolution. However, in general, $D^\alpha \varphi \notin \mathcal{D}(\mathbb{Q}_p^n)$ for $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$, where $\mathcal{D}(\mathbb{Q}_p^n)$ is the space of test functions. Consequently, the operation $D^\alpha f$ is well defined only for some distributions $f \in \mathcal{D}'(\mathbb{Q}_p^n)$. For example, D^{-1} is *defined* only on the test functions such that $\int_{\mathbb{Q}_p} \varphi(x) dx = 0$ [30, IX.2].

We recall that similar problems arise for the “ \mathbb{C} -case” fractional operators (where all functions and distributions are complex or real valued defined on spaces with real or complex coordinates): in general, the Schwartzian test function space $\mathcal{S}(\mathbb{R}^n)$ is *not invariant* under fractional operators [26], [27]. To solve this problem, in the excellent papers of P. I. Lizorkin [24], [25] a new type spaces *invariant* under fractional operators were introduced (see also [26], [27]).

Taking into account the problems mentioned above, in [3], the p -adic Lizorkin spaces of test functions and distributions were introduced, and in [3], [4] a class of pseudo-differential operators (including the Taibleson fractional operator) defined on them was constructed. The Lizorkin spaces are *invariant* under our pseudo-differential operators, and consequently, these spaces are their “natural” definition domains and can play a key role in considerations related to the fractional operators problems.

Recall that for the one-dimensional case the orthonormal complete basis of eigenfunctions (5.5) of the Vladimirov operator D^α was constructed by S. V. Kozyrev [20]. The eigenfunctions (5.5) are p -adic compactly supported wavelets. Further development and generalization of the theory of such type wavelets can be found in the papers by S. V. Kozyrev [21], [22], A. Yu. Khrennikov, and S. V. Kozyrev [16], [17], J. J. Benedetto, and R. L. Benedetto [8], and R. L. Benedetto [9].

It is typical that such type p -adic compactly supported wavelets are eigenfunctions of p -adic pseudo-differential operators. Moreover, these wavelets satisfy the condition $\int_{\mathbb{Q}_p} \varphi(x) dx = 0$ (see [20]), and, in view of Lemma 3.1, belong to the Lizorkin space $\Phi(\mathbb{Q}_p)$. In [3], there was derived the necessary and sufficient condition for multidimensional p -adic pseudo-differential operators to have such type multidimensional wavelets as eigenfunctions. Thus the wavelets theory play a key role in p -adic analysis.

Contents of the paper. In this paper problems related with the theory of multidimensional p -adic pseudo-differential operators and the theory of multidimensional p -adic wavelets are studied. Here the results of our paper [3] are intensively used.

In Sec. 2, we recall some facts from the p -adic theory of distributions [12], [28], [29], [30]. In Sec. 3, some facts from the theory of the p -adic Lizorkin

spaces [3] are recalled. In Sec. 4, we recall some facts on the multidimensional pseudo-differential operators defined in the Lizorkin space of distributions $\mathcal{D}'(\mathbb{Q}_p^n)$. The fractional Taibleson operator [28, §2], [29, III.4.] is among them. The Lizorkin spaces are *invariant* under our pseudo-differential operators. It is appropriate to mention here that the class of our operators includes the pseudo-differential operators studied in [19], [33], [34].

In Sec. 5, a *new type of p -adic compactly supported wavelets* (in one-dimensional (5.3) and multidimensional (5.17) cases) are introduced. These wavelets belong to the Lizorkin space of test functions. The Kozyrev one-dimensional wavelets [20] (see (5.5)) is a particular case of our one-dimensional wavelets (5.3). The *scaling function* of wavelets (5.3) is a characteristic function of the unit disc. The *two-scale equation* (5.7) for these wavelets is presented. However, in this paper the multiresolution analysis is not considered. The one-dimensional wavelets (5.3) and multidimensional wavelets (5.17) form orthonormal complete bases in $\mathcal{L}^2(\mathbb{Q}_p)$ and $\mathcal{L}^2(\mathbb{Q}_p^n)$, respectively (see Theorems 5.1, 5.2). Their Fourier transforms also form orthonormal complete bases in $\mathcal{L}^2(\mathbb{Q}_p)$ and $\mathcal{L}^2(\mathbb{Q}_p^n)$, respectively (see Corollary 5.1, 5.2).

In Sec. 6, the spectral theory of our pseudo-differential operators is constructed. By Theorem 6.1 the criterion (6.1) for multidimensional p -adic pseudo-differential operators (3.2) to have multidimensional wavelets (5.17) as eigenfunctions is derived. In particular, the multidimensional wavelets (5.17) are eigenfunctions of the Taibleson fractional operator (see (6.6)).

Since many p -adic models use pseudo-differential operators, in particular, fractional operator, these results on p -adic wavelets can be intensively used in applications. Moreover, p -adic wavelets can be used to construct solutions of linear and *semi-linear* pseudo-differential equations [5], [18], [23].

2. p -ADIC DISTRIBUTIONS

Now we recall some facts from the theory of p -adic distributions (generalized functions). We shall systematically use the notations and results from [30]. Let $\mathbb{N}, \mathbb{Z}, \mathbb{C}$ be the sets of positive integers, integers, complex numbers, respectively. Denote by $\mathbb{Q}_p^* = \mathbb{Q}_p \setminus \{0\}$ the multiplicative group of the field \mathbb{Q}_p . The space $\mathbb{Q}_p^n = \mathbb{Q}_p \times \cdots \times \mathbb{Q}_p$ consists of points $x = (x_1, \dots, x_n)$, where $x_j \in \mathbb{Q}_p$, $j = 1, 2, \dots, n$, $n \geq 2$. The p -adic norm on \mathbb{Q}_p^n is

$$(2.1) \quad |x|_p = \max_{1 \leq j \leq n} |x_j|_p, \quad x \in \mathbb{Q}_p^n.$$

Denote by $B_\gamma^n(a) = \{x : |x - a|_p \leq p^\gamma\}$ the ball of radius p^γ with the center at a point $a = (a_1, \dots, a_n) \in \mathbb{Q}_p^n$ and by $S_\gamma^n(a) = \{x : |x - a|_p = p^\gamma\} = B_\gamma^n(a) \setminus B_{\gamma-1}^n(a)$ its boundary (sphere), $\gamma \in \mathbb{Z}$. For $a = 0$ we set $B_\gamma^n(0) = B_\gamma^n$ and $S_\gamma^n(0) = S_\gamma^n$. For the case $n = 1$ we will omit the upper index n . Here

$$(2.2) \quad B_\gamma^n(a) = B_\gamma(a_1) \times \cdots \times B_\gamma(a_n),$$

where $B_\gamma(a_j) = \{x_j : |x_j - a_j|_p \leq p^\gamma\}$ is a disc of radius p^γ with the center at a point $a_j \in \mathbb{Q}_p$, $j = 1, 2, \dots, n$. Any two balls in \mathbb{Q}_p^n either are disjoint or one contains the other. Every point of the ball is its center.

According to [30, I.3, Examples 1, 2.], the disc B_γ is represented by the sum of $p^{\gamma-\gamma'}$ *disjoint* discs $B_{\gamma'}(a)$, $\gamma' < \gamma$:

$$(2.3) \quad B_\gamma = B_{\gamma'} \cup \bigcup_a B_{\gamma'}(a),$$

where $a = 0$ and $a = a_{-r}p^{-r} + a_{-r+1}p^{-r+1} + \dots + a_{-\gamma'+1}p^{-\gamma'+1}$ are the centers of the discs $B_{\gamma'}(a)$, $r = \gamma, \gamma-1, \gamma-2, \dots, \gamma'+1$, $0 \leq a_j \leq p-1$, $a_{-r} \neq 0$.

In particular, the disc B_0 is represented by the sum of p *disjoint* discs

$$(2.4) \quad B_0 = B_{-1} \cup \bigcup_{r=1}^{p-1} B_{-1}(r),$$

where $B_{-1}(r) = \{x \in S_0 : x_0 = r\} = r + p\mathbb{Z}_p$, $r = 1, \dots, p-1$; $B_{-1} = \{|x|_p \leq p^{-1}\} = p\mathbb{Z}_p$; and $S_0 = \{|x|_p = 1\} = \bigcup_{r=1}^{p-1} B_{-1}(r)$. Here all the discs are disjoint. We call covering (2.3), (2.4) the *canonical covering* of the disc B_0 .

A complex-valued function f defined on \mathbb{Q}_p^n is called *locally-constant* if for any $x \in \mathbb{Q}_p^n$ there exists an integer $l(x) \in \mathbb{Z}$ such that

$$f(x+y) = f(x), \quad y \in B_{l(x)}^n.$$

Let $\mathcal{E}(\mathbb{Q}_p^n)$ and $\mathcal{D}(\mathbb{Q}_p^n)$ be the linear spaces of locally-constant \mathbb{C} -valued functions on \mathbb{Q}_p^n and locally-constant \mathbb{C} -valued functions with compact supports (so-called test functions), respectively; $\mathcal{D}(\mathbb{Q}_p)$, $\mathcal{E}(\mathbb{Q}_p)$ [30, VI.1., 2.]. If $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$, according to Lemma 1 from [30, VI.1.], there exists $l \in \mathbb{Z}$, such that

$$\varphi(x+y) = \varphi(x), \quad y \in B_l^n, \quad x \in \mathbb{Q}_p^n.$$

The largest of such numbers $l = l(\varphi)$ is called the *parameter of constancy* of the function φ . Let us denote by $\mathcal{D}_N^l(\mathbb{Q}_p^n)$ the finite-dimensional space of test functions from $\mathcal{D}(\mathbb{Q}_p^n)$ having supports in the ball B_N^n and with parameters of constancy $\geq l$ [30, VI.2.]. Denote by $\mathcal{D}'(\mathbb{Q}_p^n)$ the set of all linear functionals on $\mathcal{D}(\mathbb{Q}_p^n)$ [30, VI.3.].

Let us introduce in $\mathcal{D}(\mathbb{Q}_p^n)$ a *canonical δ -sequence* $\delta_k(x) = p^{nk}\Omega(p^k|x|_p)$, and a *canonical 1-sequence* $\Delta_k(x) = \Omega(p^{-k}|x|_p)$, $k \in \mathbb{Z}$, $x \in \mathbb{Q}_p^n$, where

$$(2.5) \quad \Omega(t) = \begin{cases} 1, & 0 \leq t \leq 1, \\ 0, & t > 1. \end{cases}$$

Here $\Delta_k(x)$ is the characteristic function of the ball B_k^n . It is clear [30, VI.3., VII.1.] that $\delta_k \rightarrow \delta$, $k \rightarrow \infty$ in $\mathcal{D}'(\mathbb{Q}_p^n)$ and $\Delta_k \rightarrow 1$, $k \rightarrow \infty$ in $\mathcal{E}(\mathbb{Q}_p^n)$.

The Fourier transform of $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$ is defined by the formula

$$F[\varphi](\xi) = \int_{\mathbb{Q}_p^n} \chi_p(\xi \cdot x) \varphi(x) d^n x, \quad \xi \in \mathbb{Q}_p^n,$$

where $\chi_p(\xi \cdot x) = e^{2\pi i \sum_{j=1}^n \{\xi_j x_j\}_p}$; $\xi \cdot x$ is the scalar product of vectors; the function $\chi_p(\xi_j x_j) = e^{2\pi i \{\xi_j x_j\}_p}$ for every fixed $\xi_j \in \mathbb{Q}_p$ is an additive character

of the field \mathbb{Q}_p , $j = 1, \dots, n$; $\{x\}_p$ is the *fractional part* of a number $x \in \mathbb{Q}_p$ which is defined as follows

$$(2.6) \quad \{x\}_p = \begin{cases} 0, & \text{if } \gamma(x) \geq 0 \text{ or } x = 0, \\ p^\gamma(x_0 + x_1p + x_2p^2 + \dots + x_{|\gamma|-1}p^{|\gamma|-1}), & \text{if } \gamma(x) < 0. \end{cases}$$

The Fourier transform is a linear isomorphism $\mathcal{D}(\mathbb{Q}_p^n)$ into $\mathcal{D}(\mathbb{Q}_p^n)$. Moreover, according to [28, Lemma A.], [29, III,(3.2)], [30, VII.2.],

$$(2.7) \quad \varphi(x) \in \mathcal{D}_N^l(\mathbb{Q}_p^n) \quad \text{iff} \quad F[\varphi(x)](\xi) \in \mathcal{D}_{-l}^{-N}(\mathbb{Q}_p^n).$$

We define the Fourier transform $F[f]$ of a distribution $f \in \mathcal{D}'(\mathbb{Q}_p^n)$ by the relation [30, VII.3.]:

$$(2.8) \quad \langle F[f], \varphi \rangle = \langle f, F[\varphi] \rangle, \quad \forall \varphi \in \mathcal{D}(\mathbb{Q}_p^n).$$

Let A be a matrix and $b \in \mathbb{Q}_p^n$. Then for a distribution $f \in \mathcal{D}'(\mathbb{Q}_p^n)$ the following relation holds [30, VII,(3.3)]:

$$(2.9) \quad F[f(Ax + b)](\xi) = |\det A|_p^{-1} \chi_p(-A^{-1}b \cdot \xi) F[f(x)](A^{-1}\xi),$$

where $\det A \neq 0$. According to [30, IV,(3.1)],

$$(2.10) \quad F[\Delta_k](x) = \delta_k(x), \quad k \in \mathbb{Z}, \quad x \in \mathbb{Q}_p^n.$$

In particular, $F[\Omega(|\xi|_p)](x) = \Omega(|x|_p)$.

The convolution $f * g$ for distributions $f, g \in \mathcal{D}'(\mathbb{Q}_p^n)$ is defined (see [30, VII.1.]) as

$$(2.11) \quad \langle f * g, \varphi \rangle = \lim_{k \rightarrow \infty} \langle f(x) \times g(y), \Delta_k(x) \varphi(x + y) \rangle$$

if the limit exists for all $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$, where $f(x) \times g(y)$ is the direct product of distributions. If for distributions $f, g \in \mathcal{D}'(\mathbb{Q}_p^n)$ the convolution $f * g$ exists then [30, VII,(5.4)]

$$(2.12) \quad F[f * g] = F[f]F[g].$$

It is well known (see, e.g., [30, III.2.]) that any *multiplicative character* π of the field \mathbb{Q}_p can be represented as

$$\pi(x) \stackrel{def}{=} \pi_\alpha(x) = |x|_p^{\alpha-1} \pi_1(x), \quad x \in \mathbb{Q}_p,$$

where $\pi(p) = p^{1-\alpha}$ and $\pi_1(x)$ is a *normed multiplicative character* such that $\pi_1(x) = \pi_1(|x|_p x)$, $\pi_1(p) = \pi_1(1) = 1$, $|\pi_1(x)| = 1$. We denote $\pi_0 = |x|_p^{-1}$.

Definition 2.1. Let π_α be a multiplicative character of the field \mathbb{Q}_p .

(a) According to [1], [2], a distribution $f_m \in \mathcal{D}'(\mathbb{Q}_p)$ is said to be *associated homogeneous (in the wide sense)* of degree π_α and order m , $m \in \mathbb{N} \cup \{0\}$, if

$$\left\langle f_m, \varphi\left(\frac{x}{t}\right) \right\rangle = \pi_\alpha(t) |t|_p \langle f_m, \varphi \rangle + \sum_{j=1}^m \pi_\alpha(t) |t|_p \log_p^j |t|_p \langle f_{m-j}, \varphi \rangle$$

for all $\varphi \in \mathcal{D}(\mathbb{Q}_p)$ and $t \in \mathbb{Q}_p^*$, where $f_{m-j} \in \mathcal{D}'(\mathbb{Q}_p)$ is an associated homogeneous distribution of degree π_α and order $m-j$, $j = 1, 2, \dots, m$, i.e.,

$$f_m(tx) = \pi_\alpha(t)f_m(x) + \sum_{j=1}^m \pi_\alpha(t) \log_p^j |t|_p f_{m-j}(x), \quad t \in \mathbb{Q}_p^*.$$

If $m = 0$ we set that the above sum is empty.

(b) We say that a distribution $f \in \mathcal{D}'(\mathbb{Q}_p^n)$ is *associated homogeneous (in the wide sense)* of degree π_α and order m , $m \in \mathbb{N} \cup \{0\}$, if for all $t \in \mathbb{Q}_p^*$ we have

$$f_m(tx) = f_m(tx_1, \dots, tx_n) = \pi_\alpha(t)f_m(x) + \sum_{j=1}^m \pi_\alpha(t) \log_p^j |t|_p f_{m-j}(x),$$

where $x = (x_1, \dots, x_n) \in \mathbb{Q}_p^n$, $f_{m-j} \in \mathcal{D}'(\mathbb{Q}_p^n)$ is an associated homogeneous distribution of degree π_α and order $m-j$, $j = 1, 2, \dots, m$. An *associated homogeneous (in the wide sense)* distribution of degree $\pi_\alpha(t) = |t|_p^{\alpha-1}$ and order m is called *associated homogeneous* of degree $\alpha-1$ and order m .

(c) An associated homogeneous distribution (in the wide sense) of order $m = 1$ is called *associated homogeneous* distribution (see [11] and [1], [2]).

(d) An associated homogeneous distribution of degree π_α and order $m = 0$ is called *homogeneous* distribution of degree π_α , i.e.,

$$f_0(tx) = f_0(tx_1, \dots, tx_n) = \pi_\alpha(t)f_0(x), \quad x = (x_1, \dots, x_n) \in \mathbb{Q}_p^n.$$

(for one-dimensional case see [12, Ch.II, §2.3.], [30, VIII.1.]).

The multidimensional homogeneous distribution $|x|_p^{\alpha-n} \in \mathcal{D}'(\mathbb{Q}_p^n)$ of degree $\alpha-n$ is constructed as follows. If $\operatorname{Re} \alpha > 0$ then the function $|x|_p^{\alpha-n}$ generates a regular functional

$$(2.13) \quad \langle |x|_p^{\alpha-n}, \varphi \rangle = \int_{\mathbb{Q}_p^n} |x|_p^{\alpha-n} \varphi(x) d^n x, \quad \forall \varphi \in \mathcal{D}(\mathbb{Q}_p^n),$$

where $|x|_p$, $x \in \mathbb{Q}_p^n$ is given by (2.1). If $\operatorname{Re} \alpha \leq 0$ this distribution is defined by means of analytic continuation [28, (*)], [29, III, (4.3)], [30, VIII, (4.2)]:

$$(2.14) \quad \begin{aligned} \langle |x|_p^{\alpha-n}, \varphi \rangle &= \int_{B_0^n} |x|_p^{\alpha-n} (\varphi(x) - \varphi(0)) d^n x \\ &+ \int_{\mathbb{Q}_p^n \setminus B_0^n} |x|_p^{\alpha-n} \varphi(x) d^n x + \varphi(0) \frac{1 - p^{-n}}{1 - p^{-\alpha}}, \end{aligned}$$

for all $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$, $\alpha \neq \mu_j = \frac{2\pi i}{\ln p} j$, $j \in \mathbb{Z}$. The distribution $|x|_p^{\alpha-n}$ is an entire function of the complex variable α everywhere except the points μ_j , $j \in \mathbb{Z}$, where it has simple poles with residues $\frac{1-p^{-n}}{\log p} \delta(x)$.

Similarly to the one-dimensional case [1], [2], one can construct the distribution $P(\frac{1}{|x|_p^n})$ called the principal value of the function $\frac{1}{|x|_p^n}$, $x \in \mathbb{Q}_p^n$:

$$(2.15) \quad \left\langle P\left(\frac{1}{|x|_p^n}\right), \varphi \right\rangle = \int_{B_0^n} \frac{\varphi(x) - \varphi(0)}{|x|_p^n} d^n x + \int_{\mathbb{Q}_p^n \setminus B_0^n} \frac{\varphi(x)}{|x|_p^n} d^n x,$$

for all $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$. It is easy to show that this distribution is *associated homogeneous* of degree $-n$ and order 1 (see [1], [2]).

The Fourier transform of $|x|_p^{\alpha-n}$ is given by the formula from [28, Theorem 2.], [29, III, Theorem (4.5)], [30, VIII, (4.3)]

$$(2.16) \quad F[|x|_p^{\alpha-n}] = \Gamma_p^{(n)}(\alpha) |\xi|_p^{-\alpha}, \quad \alpha \neq 0, n$$

where the n -dimensional Γ -function $\Gamma_p^{(n)}(\alpha)$ is given by the following formulas (see [28, Theorem 1.], [29, III, Theorem (4.2)], [30, VIII, (4.4)]):

$$(2.17) \quad \begin{aligned} \Gamma_p^{(n)}(\alpha) &\stackrel{\text{def}}{=} \lim_{k \rightarrow \infty} \int_{p^{-k} \leq |x|_p \leq p^k} |x|_p^{\alpha-n} \chi_p(u \cdot x) d^n x \\ &= \int_{\mathbb{Q}_p^n} |x|_p^{\alpha-n} \chi_p(x_1) d^n x = \frac{1 - p^{\alpha-n}}{1 - p^{-\alpha}} \end{aligned}$$

where $|u|_p = 1$, and the last integrals in the right-hand side of (2.17) are defined by means of analytic continuation with respect to the parameter α . Here $\Gamma_p^{(1)}(\alpha) = \Gamma_p(\alpha) = \int_{\mathbb{Q}_p} |x|_p^{\alpha-1} \chi_p(x) dx = \frac{1-p^{\alpha-1}}{1-p^{-\alpha}}$.

3. THE p -ADIC LIZORKIN SPACES

Let us introduce the p -adic *Lizorkin space of test functions*

$$\Phi(\mathbb{Q}_p^n) = \{\phi : \phi = F[\psi], \psi \in \Psi(\mathbb{Q}_p^n)\},$$

where

$$\Psi(\mathbb{Q}_p^n) = \{\psi(\xi) \in \mathcal{D}(\mathbb{Q}_p^n) : \psi(0) = 0\}.$$

Here $\Psi(\mathbb{Q}_p^n), \Phi(\mathbb{Q}_p^n) \subset \mathcal{D}(\mathbb{Q}_p^n)$. The space $\Phi(\mathbb{Q}_p^n)$ can be equipped with the topology of the space $\mathcal{D}(\mathbb{Q}_p^n)$ which makes $\Phi(\mathbb{Q}_p^n)$ a complete space.

In view of (2.7), the following lemma holds.

Lemma 3.1. ([3], [4]) (a) $\phi \in \Phi(\mathbb{Q}_p^n)$ iff $\phi \in \mathcal{D}(\mathbb{Q}_p^n)$ and

$$(3.1) \quad \int_{\mathbb{Q}_p^n} \phi(x) d^n x = 0.$$

(b) $\phi \in \mathcal{D}_N^l(\mathbb{Q}_p^n) \cap \Phi(\mathbb{Q}_p^n)$, i.e., $\int_{B_N^n} \phi(x) d^n x = 0$, iff $\psi = F^{-1}[\phi] \in \mathcal{D}_{-l}^{-N}(\mathbb{Q}_p^n) \cap \Psi(\mathbb{Q}_p^n)$, i.e., $\psi(\xi) = 0$, $\xi \in B_{-N}^n$.

In fact, for $n = 1$, this lemma was proved in [30, IX.2.]. Unlike the classical Lizorkin space, any function $\psi(\xi) \in \Phi(\mathbb{Q}_p^n)$ is equal to zero not only at $\xi = 0$ but in a ball $B^n \ni 0$, as well.

Let $\Phi'(\mathbb{Q}_p^n)$ denote the topological dual of the space $\Phi(\mathbb{Q}_p^n)$. We call it the p -adic *Lizorkin space of distributions*.

By Ψ^\perp and Φ^\perp we denote the subspaces of functionals in $\mathcal{D}'(\mathbb{Q}_p^n)$ orthogonal to $\Psi(\mathbb{Q}_p^n)$ and $\Phi(\mathbb{Q}_p^n)$, respectively. Thus $\Psi^\perp = \{f \in \mathcal{D}'(\mathbb{Q}_p^n) : f = C\delta, C \in \mathbb{C}\}$ and $\Phi^\perp = \{f \in \mathcal{D}'(\mathbb{Q}_p^n) : f = C, C \in \mathbb{C}\}$.

Proposition 3.1. ([3])

$$\Phi'(\mathbb{Q}_p^n) = \mathcal{D}'(\mathbb{Q}_p^n)/\Phi^\perp, \quad \Psi'(\mathbb{Q}_p^n) = \mathcal{D}'(\mathbb{Q}_p^n)/\Psi^\perp.$$

The space $\Phi'(\mathbb{Q}_p^n)$ can be obtained from $\mathcal{D}'(\mathbb{Q}_p^n)$ by “sifting out” constants. Thus two distributions in $\mathcal{D}'(\mathbb{Q}_p^n)$ differing by a constant are indistinguishable as elements of $\Phi'(\mathbb{Q}_p^n)$.

Similarly to (2.8), we define the Fourier transforms of distributions $f \in \Phi'_\times(\mathbb{Q}_p^n)$ and $g \in \Psi'_\times(\mathbb{Q}_p^n)$ by the relations:

$$(3.2) \quad \begin{aligned} \langle F[f], \psi \rangle &= \langle f, F[\psi] \rangle, & \forall \psi \in \Psi(\mathbb{Q}_p^n), \\ \langle F[g], \phi \rangle &= \langle g, F[\phi] \rangle, & \forall \phi \in \Phi(\mathbb{Q}_p^n). \end{aligned}$$

By definition, $F[\Phi(\mathbb{Q}_p^n)] = \Psi(\mathbb{Q}_p^n)$ and $F[\Psi(\mathbb{Q}_p^n)] = \Phi(\mathbb{Q}_p^n)$, i.e., (3.2) give well defined objects.

4. PSEUDO-DIFFERENTIAL OPERATORS IN THE LIZORKIN SPACE

4.1. Pseudo-differential operators. Consider a class of pseudo-differential operators in the Lizorkin space of the test functions $\Phi(\mathbb{Q}_p^n)$

$$(4.1) \quad \begin{aligned} (A\phi)(x) &= F^{-1}[\mathcal{A}(\xi) F[\phi](\xi)](x) \\ &= \int_{\mathbb{Q}_p^n} \int_{\mathbb{Q}_p^n} \chi_p((y-x) \cdot \xi) \mathcal{A}(\xi) \phi(y) d^n \xi d^n y, \quad \phi \in \Phi(\mathbb{Q}_p^n) \end{aligned}$$

with symbols $\mathcal{A}(\xi) \in \mathcal{E}(\mathbb{Q}_p^n \setminus \{0\})$.

Lemma 4.1. *The Lizorkin space $\Phi(\mathbb{Q}_p^n)$ is invariant under the pseudo-differential operators (4.1). Moreover, $A(\Phi(\mathbb{Q}_p^n)) = \Phi(\mathbb{Q}_p^n)$.*

Proof. In view of (2.7) and results of Sec. 3, functions $F[\phi](\xi)$ and $\mathcal{A}(\xi)F[\phi](\xi)$ belong to $\Psi(\mathbb{Q}_p^n)$, and, consequently, $(A\phi)(x) \in \Phi(\mathbb{Q}_p^n)$, i.e., $A(\Phi(\mathbb{Q}_p^n)) \subset \Phi(\mathbb{Q}_p^n)$. Thus the pseudo-differential operators (4.1) are well defined, and the Lizorkin space $\Phi(\mathbb{Q}_p^n)$ is invariant under them. Moreover, any function from $\Psi(\mathbb{Q}_p^n)$ can be represented as $\psi(\xi) = \mathcal{A}(\xi)\psi_1(\xi)$, $\psi_1 \in \Psi(\mathbb{Q}_p^n)$. This implies that $A(\Phi(\mathbb{Q}_p^n)) = \Phi(\mathbb{Q}_p^n)$. \square

If we define a conjugate pseudo-differential operator A^T as

$$(4.2) \quad (A^T \phi)(x) = F^{-1}[\mathcal{A}(-\xi) F[\phi](\xi)](x) = \int_{\mathbb{Q}_p^n} \chi_p(-x \cdot \xi) \mathcal{A}(-\xi) F[\phi](\xi) d^n \xi$$

then one can define the operator A in the Lizorkin space of distributions: for $f \in \Phi'(\mathbb{Q}_p^n)$ we have

$$(4.3) \quad \langle Af, \phi \rangle = \langle f, A^T \phi \rangle, \quad \forall \phi \in \Phi(\mathbb{Q}_p^n).$$

It is clear that

$$(4.4) \quad Af = F^{-1}[\mathcal{A}F[f]] \in \Phi'(\mathbb{Q}_p^n),$$

i.e., the Lizorkin space of distributions $\Phi'(\mathbb{Q}_p^n)$ is invariant under pseudo-differential operators A . Moreover, in view of Lemma 4.1, $A(\Phi'(\mathbb{Q}_p^n)) = \Phi'(\mathbb{Q}_p^n)$.

If A, B are pseudo-differential operators with symbols $\mathcal{A}(\xi), \mathcal{B}(\xi) \in \mathcal{E}(\mathbb{Q}_p^n \setminus \{0\})$, respectively, then the operator AB is well defined and represented by the formula

$$(AB)f = F^{-1}[\mathcal{A}\mathcal{B}F[f]] \in \Phi'(\mathbb{Q}_p^n).$$

If $\mathcal{A}(\xi) \neq 0, \xi \in \mathbb{Q}_p^n \setminus \{0\}$ then we define the inverse pseudo-differential operator by the formula

$$A^{-1}f = F^{-1}[\mathcal{A}^{-1}F[f]], \quad f \in \Phi'(\mathbb{Q}_p^n).$$

Thus the family of pseudo-differential operators A with symbols $\mathcal{A}(\xi) \neq 0, \xi \in \mathbb{Q}_p^n \setminus \{0\}$ forms an Abelian group.

If the symbol $\mathcal{A}(\xi)$ of the operator A is a *homogeneous* or an *associated homogeneous* function (see Definition 2.1) then the pseudo-differential operator A is called *homogeneous* or *associated homogeneous*.

4.2. The Taibleson fractional operator. Let us consider a pseudo-differential operator D_x^α with the symbol $\mathcal{A}(\xi) = |\xi|_p^\alpha$. Thus, according to (4.1),

$$(4.5) \quad (D_x^\alpha \phi)(x) = F^{-1}[\xi^\alpha F[\phi](\xi)](x), \quad \phi \in \Phi(\mathbb{Q}_p^n).$$

This multi-dimensional Taibleson fractional operator was introduced in [28, §2], [29, III.4.] on the space of distributions $\mathcal{D}'(\mathbb{Q}_p^n)$ for $\alpha \in \mathbb{C}, \alpha \neq -n$.

In view of formulas (2.12), (2.16), (2.17), the relation (4.5) can be rewritten as a convolution

$$(D_x^\alpha \phi)(x) \stackrel{def}{=} \kappa_{-\alpha}(x) * \phi(x) = \langle \kappa_{-\alpha}(x), \phi(x - \xi) \rangle, \quad x \in \mathbb{Q}_p^n,$$

where $\phi \in \Phi(\mathbb{Q}_p^n), \alpha \neq 0, -n$. Here the distribution from $\mathcal{D}'(\mathbb{Q}_p^n)$

$$(4.6) \quad \kappa_\alpha(x) = \frac{|x|_p^{\alpha-n}}{\Gamma_p^{(n)}(\alpha)}, \quad \alpha \neq 0, n, \quad x \in \mathbb{Q}_p^n,$$

is called the multidimensional *Riesz kernel* [28, §2], [29, III.4.], where the function $|x|_p, x \in \mathbb{Q}_p^n$ is given by (2.1). The Riesz kernel has a removable singularity at $\alpha = 0$ and according to [28, §2], [29, III.4.], [30, VIII.2], we obtain that $\langle \kappa_0(x), \varphi(x) \rangle \stackrel{def}{=} \lim_{\alpha \rightarrow 0} \langle \kappa_\alpha(x), \varphi(x) \rangle = \varphi(0)$, for all $\varphi \in \mathcal{D}(\mathbb{Q}_p^n)$, i.e.,

$$(4.7) \quad \kappa_0(x) \stackrel{def}{=} \lim_{\alpha \rightarrow 0} \kappa_\alpha(x) = \delta(x).$$

Using (2.13), (2.17), (4.6), and taking into account (3.1) (for details, see [3], [4]), we define $\kappa_n(\cdot)$ as a distribution from the *Lizorkin space of distributions* $\Phi'(\mathbb{Q}_p^n)$:

$$(4.8) \quad \kappa_n(x) \stackrel{\text{def}}{=} \lim_{\alpha \rightarrow n} \kappa_\alpha(x) = -\frac{1-p^{-n}}{\log p} \log |x|_p.$$

With the help of (2.16), (4.7), (4.8), it is easy to see that

$$(4.9) \quad \kappa_\alpha(x) * \kappa_\beta(x) = \kappa_{\alpha+\beta}(x), \quad \alpha, \beta \in \mathbb{C},$$

holds in the sense of the Lizorkin space $\Phi'(\mathbb{Q}_p^n)$.

In view of (4.7), (4.8), the multi-dimensional Taibleson operator on the Lizorkin space of test functions is defined for all $\alpha \in \mathbb{C}$ as

$$(4.10) \quad (D_x^\alpha \phi)(x) \stackrel{\text{def}}{=} \kappa_{-\alpha}(x) * \phi(x) = \langle \kappa_{-\alpha}(x), \phi(x - \xi) \rangle, \quad x \in \mathbb{Q}_p^n,$$

where $\phi \in \Phi(\mathbb{Q}_p^n)$.

If $\alpha \neq n$ then the Riesz kernel $\kappa_\alpha(x)$ is a *homogeneous* distribution of degree $\alpha - n$, and if $\alpha = n$ then the Riesz kernel is an *associated homogeneous* distribution of degree 0 and order 1 (see Definitions 2.1,(b),(d)). Thus the Taibleson fractional operator D_x^α , $\alpha \neq -n$ is a *homogeneous* pseudo-differential operator of degree α , and D_x^{-n} is an *associated homogeneous* pseudo-differential operator of degree $-n$ and order 1 with the symbol $\mathcal{A}(\xi) = P(|\xi|_p^{-n})$ (see (2.15)).

According to Lemma 4.1, the Lizorkin space $\Phi(\mathbb{Q}_p^n)$ is invariant under the Taibleson fractional operator D_x^α and $D_x^\alpha(\Phi(\mathbb{Q}_p^n)) = \Phi(\mathbb{Q}_p^n)$ [3].

In view of (4.2), (4.3), $(D_x^\alpha)^T = D_x^\alpha$ and for $f \in \Phi'(\mathbb{Q}_p^n)$ we have

$$(4.11) \quad \langle D_x^\alpha f, \phi \rangle \stackrel{\text{def}}{=} \langle f, D_x^\alpha \phi \rangle, \quad \forall \phi \in \Phi(\mathbb{Q}_p^n).$$

It is clear that $D_x^\alpha(\Phi'(\mathbb{Q}_p^n)) = \Phi'(\mathbb{Q}_p^n)$. Moreover, in view of (4.9), the family of operators D_x^α , $\alpha \in \mathbb{C}$ on the Lizorkin space forms an Abelian group: if $f \in \Phi'(\mathbb{Q}_p^n)$ then $D_x^\alpha D_x^\beta f = D_x^\beta D_x^\alpha f = D_x^{\alpha+\beta} f$, $D_x^\alpha D_x^{-\alpha} f = f$, $\alpha, \beta \in \mathbb{C}$.

5. p -ADIC WAVELETS

5.1. One-dimensional p -adic wavelets. Let $n = 1$. Consider the set

$$I_p = \{a = p^{-\gamma}(a_0 + a_1 p + \cdots + a_{\gamma-1} p^{\gamma-1}) :$$

$$(5.1) \quad \gamma \in \mathbb{N}; a_j = 0, 1, \dots, p-1; j = 0, 1, \dots, \gamma-1\}.$$

This set can be identified with the factor group $\mathbb{Q}_p/\mathbb{Z}_p$. Let

$$J_{p;m} = \{s = p^{-m}(s_0 + s_1 p + \cdots + s_{m-1} p^{m-1}) :$$

$$(5.2) \quad s_j = 0, 1, \dots, p-1; j = 0, 1, \dots, m-1; s_0 \neq 0\},$$

where $m \geq 1$ is a *fixed* positive integer.

Let us introduce the function $\theta_s^{(m)}(x) = \chi_p(sx)\Omega(|x|_p)$, $x \in \mathbb{Q}_p$, $s \in J_{p;m}$, and the functions generated by its dilatations and translations:

$$(5.3) \quad \theta_{\gamma sa}^{(m)}(x) = p^{-\gamma/2} \chi_p(s(p^\gamma x - a)) \Omega(|p^\gamma x - a|_p), \quad x \in \mathbb{Q}_p,$$

where $\gamma \in \mathbb{Z}$, $s \in J_{p;m}$, $a \in I_p$, $\Omega(t)$ is the characteristic function (2.5) of the segment $[0, 1]$.

Making the change of variables $\xi = p^\gamma x - a$ and taking into account (2.10), we obtain

$$(5.4) \quad \int_{\mathbb{Q}_p} \theta_{\gamma sa}^{(m)}(x) dx = p^{\gamma/2} \int_{\mathbb{Q}_p} \chi_p(s\xi) \Omega(|\xi|_p) d\xi = p^{\gamma/2} \Omega(|s|_p) = 0.$$

Thus, in view of Theorem 5.1 (see below), one can see that the functions (5.3) are *p-adic wavelets*. Moreover, according to (5.4) and Lemma 3.1, the functions $\theta_{\gamma sa}^{(m)}(x)$ belong to the Lizorkin space $\Phi(\mathbb{Q}_p)$.

It is clear that for any $\gamma \in \mathbb{Z}$ and $s \in J_{p;m}$ the functions (5.4) are *periodical* with the periods $T_{\gamma s} \in p^{m-\gamma}\mathbb{Z}_p$.

In the case $m = 1$, i.e., for $s = p^{-1}j$, $j = 1, 2, \dots, p-1$ these wavelets coincide with the Kozyrev wavelets [20]:

$$(5.5) \quad \theta_{\gamma sa}^{(1)}(x) = \theta_{\gamma ja}(x) = p^{-\gamma/2} \chi_p(p^{-1}j(p^\gamma x - a)) \Omega(|p^\gamma x - a|_p), \quad x \in \mathbb{Q}_p,$$

$\gamma \in \mathbb{Z}$, $j = 1, 2, \dots, p-1$, $a \in I_p$.

In particular, $\theta_s^{(1)}(x) = \theta_j(x) = \chi_p(p^{-1}jx)\Omega(|x|_p)$ for $j = 1$. Since $|x|_p \leq 1$, $x \in \mathbb{Q}_p$, i.e., $x = x_0 + x_1p + x_2p^2 + \dots$, we have $p^{-1}x = p^{-1}x_0 + x_1 + x_2p + \dots$, i.e., the fractional part (2.6) of a number $p^{-1}x$ is equal to $\{p^{-1}x\}_p = p^{-1}x_0$. According to (2.4),

$$(5.6) \quad \theta_1(x) = \chi_p(p^{-1}x)\Omega(|x|_p) = \begin{cases} 0, & |x|_p \geq p, \\ e^{2\pi i \frac{x}{p}}, & x \in B_{-1}(r), r = 1, \dots, p-1, \\ 1, & x \in B_{-1}. \end{cases}$$

Thus the function $\theta_1(x) = \chi_p(p^{-1}x)\Omega(|x|_p)$ takes values in the set $\{0, e^{2\pi i \frac{x}{p}} : r = 0, 1, \dots, p-1\}$ of $p+1$ elements.

Now we consider $\theta_s^{(1)}(x) = \theta_j(x) = \chi_p(p^{-1}jx)\Omega(|x|_p)$. Since $|jx|_p \leq 1$, $x \in \mathbb{Q}_p$, we have $jx = y_0 + y_1p + y_2p^2 + \dots$, $p^{-1}jx = p^{-1}y_0 + y_1 + y_2p + \dots$, and $\{p^{-1}jx\}_p = p^{-1}y_0$ (see (2.6)). Thus,

$$\theta_j(x) = \chi_p(p^{-1}jx)\Omega(|x|_p) = \begin{cases} 0, & |x|_p \geq p, \\ e^{2\pi i \{\frac{jx}{p}\}_p}, & x \in B_{-1}(r), r = 1, \dots, p-1, \\ 1, & x \in B_{-1}. \end{cases}$$

It is clear that for the Kozyrev wavelets the *scaling function* is the characteristic function of the unit disc $\Delta_0(x) = \Omega(|x|_p)$, $x \in \mathbb{Q}_p$, and in view of (2.4) it satisfies the *two-scale equation*:

$$(5.7) \quad \Delta_0(x) = p^{-1/2} \sum_{r=0}^{p-1} h_r \Delta_0\left(\frac{1}{p}x - \frac{r}{p}\right), \quad x \in \mathbb{Q}_p,$$

where $h_r = p^{1/2}$. Relations (5.6), (5.7) imply that

$$(5.8) \quad \theta_1(x) = \chi_p(p^{-1}x)\Omega(|x|_p) = p^{-1/2} \sum_{r=0}^{p-1} \tilde{h}_r \Delta_0\left(\frac{1}{p}x - \frac{r}{p}\right), \quad x \in \mathbb{Q}_p,$$

where $\tilde{h}_r = p^{1/2} e^{2\pi i \frac{r}{p}}$, $r = 0, 1, \dots, p-1$. Similarly to (5.8), we have

$$(5.9) \quad \theta_j(x) = \chi_p(p^{-1}jx)\Omega(|x|_p) = p^{-1/2} \sum_{r=0}^{p-1} \tilde{h}_r \Delta_0\left(\frac{1}{p}x - \frac{r}{p}\right), \quad x \in \mathbb{Q}_p,$$

where $\tilde{h}_r = p^{1/2} e^{2\pi i \{\frac{jr}{p}\}_p}$, $r = 0, 1, \dots, p-1$.

In the same way we consider the function $\theta_s^{(m)}(x) = \chi_p(sx)\Omega(|x|_p)$, $s \in J_{p;m}$. Let $B_0 = \cup_a B_{-m}(a) \cup B_{-m}$ be the *canonical covering* (2.3) of the disc B_0 with p^m discs, $m \geq 1$, where $a = 0$ and $a = a_r p^r + a_{r+1} p^{r+1} + \dots + a_{m-1} p^{m-1}$ is the center of the discs B_{-m} and $B_{-m}(a)$, respectively, $r = 0, 1, 2, \dots, m-1$, $0 \leq a_j \leq p-1$, $a_r \neq 0$.

For $x \in B_{-m}(a)$, $s \in J_{p;m}$, we have $x = a + p^m(y_0 + y_1 p + y_2 p^2 + \dots)$, $s = p^{-m}(s_0 + s_1 p + \dots + s_{m-1} p^{m-1})$, $s_0 \neq 0$; $sx = sa + \xi$, $\xi \in \mathbb{Z}_p$; and $\{sx\}_p = \{sa\}_p = \{p^{r-m}(a_r + a_{r+1} p + \dots + a_{m-1} p^{m-r-1})(s_0 + s_1 p + \dots + s_{m-1} p^{m-1})\}_p$, $r = 0, 1, 2, \dots, m-1$, (see (2.6)). Thus,

$$\begin{aligned} \theta_s^{(m)}(x) &= \chi_p(sx)\Omega(|x|_p) \\ &= \begin{cases} 0, & |x|_p \geq p, \\ e^{2\pi i \{sa\}_p}, & x \in B_{-m}(a), \quad a = \sum_{l=r}^{m-1} a_l p^l, \\ 1, & x \in B_{-m}, \end{cases} \end{aligned}$$

where $s = p^{-m}(s_0 + s_1 p + \dots + s_{m-1} p^{m-1})$, $0 \leq s_j \leq p-1$, $j = 0, 1, \dots, m-1$, $s_0 \neq 0$; $a = a_r p^r + a_{r+1} p^{r+1} + \dots + a_{m-1} p^{m-1}$, $0 \leq a_j \leq p-1$, $a_r \neq 0$, $r = 0, 1, \dots, m-1$. Thus the function $\theta_s^{(m)}(x) = \chi_p(sx)\Omega(|x|_p)$ takes values in the set $\{0, 1, e^{2\pi i \{sa\}_p}\}$ of $p^m + 1$ elements.

In this case, using the *scaling function*, we obtain

$$(5.10) \quad \theta_s^{(m)}(x) = \chi_p(sx)\Omega(|x|_p) = p^{-m/2} \sum_a \tilde{h}_a \Delta_0\left(\frac{1}{p^m}x - \frac{a}{p^m}\right),$$

$x \in \mathbb{Q}_p$, where $\tilde{h}_0 = p^{m/2}$; $\tilde{h}_a = p^{m/2} e^{2\pi i \{sa\}_p}$, $a = a_r p^r + a_{r+1} p^{r+1} + \dots + a_{m-1} p^{m-1}$, $r = 0, 1, \dots, m-1$, $0 \leq a_j \leq p-1$, $a_r \neq 0$.

Theorem 5.1. *The functions (5.3) form an orthonormal complete basis in $\mathcal{L}^2(\mathbb{Q}_p)$ (p -adic wavelet basis).*

Proof. Consider the scalar product

$$\begin{aligned} &(\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = p^{-(\gamma + \gamma')/2} \\ (5.11) \quad &\times \int_{\mathbb{Q}_p} \chi_p(s'(p^{\gamma'} x - a') - s(p^\gamma x - a)) \Omega(|p^\gamma x - a|_p) \Omega(|p^{\gamma'} x - a'|_p) dx. \end{aligned}$$

If $\gamma \leq \gamma'$, according to formula [30, VII.1], [20]

$$(5.12) \quad \Omega(|p^\gamma x - a|_p) \Omega(|p^{\gamma'} x - a'|_p) = \Omega(|p^\gamma x - a|_p) \Omega(|p^{\gamma'-\gamma} a - a'|_p),$$

(5.11) can be rewritten as

$$(5.13) \quad \begin{aligned} & (\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = p^{-(\gamma+\gamma')/2} \Omega(|p^{\gamma'-\gamma} a - a'|_p) \\ & \times \int_{\mathbb{Q}_p} \chi_p(s'(p^{\gamma'} x - a') - s(p^\gamma x - a)) \Omega(|p^\gamma x - a|_p) dx. \end{aligned}$$

Let $\gamma < \gamma'$. Making the change of variables $\xi = p^\gamma x - a$ and taking into account (2.10), from (5.13) we obtain

$$(5.14) \quad \begin{aligned} & (\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = p^{-(\gamma+\gamma')/2} \chi_p(s'(p^{\gamma'-\gamma} a - a')) \\ & \times \Omega(|p^{\gamma'-\gamma} a - a'|_p) \int_{\mathbb{Q}_p} \chi_p((p^{\gamma'-\gamma} s' - s)\xi) \Omega(|\xi|_p) d\xi \\ & = p^{-(\gamma+\gamma')/2} \chi_p(s'(p^{\gamma'-\gamma} a - a')) \Omega(|p^{\gamma'-\gamma} a - a'|_p) \Omega(|p^{\gamma'-\gamma} s' - s|_p). \end{aligned}$$

Since

$$\begin{aligned} p^{\gamma'-\gamma} s' &= p^{\gamma'-\gamma-m} (s'_0 + s'_1 p + \cdots + s'_{\gamma-1} p^{m-1}), \\ s &= p^{-m} (s_0 + s_1 p + \cdots + s_{\gamma-1} p^{m-1}), \end{aligned}$$

where $s'_0, s_0 \neq 0$, $\gamma' - \gamma \leq 1$, it is clear that fractional part $\{p^{\gamma'-\gamma} s' - s\}_p \neq 0$. Thus $\Omega(|p^{\gamma'-\gamma} s' - s|_p) = 0$ and $(\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = 0$.

Consequently, the scalar product $(\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = 0$ can be nonzero only if $\gamma = \gamma'$. In this case (5.14) implies

$$(5.15) \quad (\theta_{\gamma s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = p^{-\gamma} \chi_p(s'(a - a')) \Omega(|a - a'|_p) \Omega(|s' - s|_p),$$

where $\Omega(|a - a'|_p) = \delta_{a'a}$, $\Omega(|s' - s|_p) = \delta_{s's}$, and $\delta_{s's}$, $\delta_{a'a}$ are the Kronecker symbols.

Since $\int_{\mathbb{Q}_p} \Omega(|p^\gamma x - a|_p) dx = p^\gamma$ [30, IV,(2.3)], formulas (5.14), (5.15) imply that

$$(5.16) \quad (\theta_{\gamma' s' a'}^{(m)}(x), \theta_{\gamma s a}^{(m)}(x)) = \delta_{\gamma'\gamma} \delta_{s's} \delta_{a'a}.$$

Thus the system of functions (5.3) is orthonormal.

To prove the completeness of the system of functions (5.3), we repeat the corresponding proof [20] almost word for word. Recall that the system of the characteristic functions of the discs B_k is complete in $\mathcal{L}^2(\mathbb{Q}_p)$. Consequently, taking into account that the system of functions $\{\theta_{\gamma s a}^{(m)}(x) : \gamma \in \mathbb{Z}, s \in J_{p;m}, a \in I_p\}$ is invariant under dilatations and translations, in order to prove that it is a complete system, it is sufficient to verify the Parseval identity for the characteristic function $\Omega(|x|_p)$.

If $0 \leq \gamma$, according to (5.12), (2.10),

$$(\Omega(|x|_p), \theta_{\gamma s a}^{(m)}(x)) = p^{-\gamma/2} \Omega(|-a|_p) \int_{\mathbb{Q}_p} \chi_p(s(p^\gamma x - a)) \Omega(|x|_p) dx$$

$$= p^{-\gamma/2} \chi_p(-sp^\gamma a) \Omega(|sp^\gamma|_p) \Omega(|-a|_p) = \begin{cases} 0, & a \neq 0, \\ 0, & a = 0, \gamma \leq m-1, \\ p^{-\gamma/2}, & a = 0, \gamma \geq m. \end{cases}$$

If $0 > \gamma$, according to (5.12), (2.10),

$$\begin{aligned} (\Omega(|x|_p), \theta_{\gamma sa}^{(m)}(x)) &= p^{-\gamma/2} \Omega(|p^{-\gamma}a|_p) \int_{\mathbb{Q}_p} \chi_p(s(p^\gamma x - a)) \Omega(|p^\gamma x - a|_p) dx \\ &= p^{-\gamma/2} \Omega(|p^{-\gamma}a|_p) \int_{\mathbb{Q}_p} \chi_p(s\xi) \Omega(|\xi|_p) d\xi = p^{-\gamma/2} \Omega(|p^{-\gamma}a|_p) \Omega(|s|_p) = 0. \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{\gamma \in \mathbb{Z}, s \in J_{p;m}, a \in I_p} |(\Omega(|x|_p), \theta_{\gamma sa}^{(m)}(x))|^2 &= \sum_{\gamma=m}^{\infty} \sum_{s \in J_{p;m}} p^{-\gamma} \\ &= p^{m-1} (p-1) \frac{p^{-m}}{1-p^{-1}} = 1 = |(\Omega(|x|_p), \Omega(|x|_p))|^2. \end{aligned}$$

Thus the system of functions (5.3) is an orthonormal basis in $\mathcal{L}^2(\mathbb{Q}_p)$ (p -adic wavelet basis). \square

Corollary 5.1. *The functions*

$$\tilde{\theta}_{\gamma sa}^{(m)} = F[\theta_{\gamma sa}^{(m)}](\xi) = p^{\gamma/2} \chi_p(p^{-\gamma}a \cdot \xi) \Omega(|s + p^{-\gamma}\xi|_p), \quad \xi \in \mathbb{Q}_p,$$

form an orthonormal complete basis in $\mathcal{L}^2(\mathbb{Q}_p)$, $a \in I_p$; $s \in J_{p;m}$; $m \geq 1$ is a fixed positive integer.

The proof follows from Theorem 5.1, formula (6.3) (see below) and the Parseval formula [30, VII,(4.1)]

5.2. Multidimensional p -adic wavelets. Let us introduce n -dimensional functions generated by the n -direct product of the one-dimensional p -adic wavelets (5.3):

$$(5.17) \quad \Theta_{\gamma sa}^{(m)}(x) = p^{-n\gamma/2} \chi_p(s \cdot (p^\gamma x - a)) \Omega(|p^\gamma x - a|_p),$$

$x = (x_1, \dots, x_n) \in \mathbb{Q}_p^n$, where $\gamma \in \mathbb{Z}$; $a = (a_1, \dots, a_n) \in I_p^n$; $s = (s_1, \dots, s_n) \in J_{p;m}^n$; $m = (m_1, \dots, m_n)$, $m_j \geq 1$ is a fixed positive integer, $j = 1, 2, \dots, n$. Here $I_p^n = I_p \times \dots \times I_p$ and $J_{p;m}^n = J_{p;m_1} \times \dots \times J_{p;m_n}$ are the n -direct products of the corresponding sets (5.1) and (5.2).

Using (5.4), (2.2), it is easy to verify that

$$(5.18) \quad \int_{\mathbb{Q}_p^n} \Theta_{\gamma sa}^{(m)}(x) d^n x = 0.$$

Thus the functions (5.17) are n -dimensional p -adic wavelets. According to (5.18) and Lemma 3.1, $\Theta_{\gamma sa}^{(m)}(x)$ belong to the Lizorkin space $\Phi(\mathbb{Q}_p^n)$.

For any $\gamma \in \mathbb{Z}$ and $s = (s_1, \dots, s_n) \in J_{p;m}^n$ the functions (5.17) are periodical with the vector periods $T_{\gamma s} = (T_{1|\gamma s}, \dots, T_{n|\gamma s}) \in p^{m-\gamma} \mathbb{Z}_p^n$.

In view of (2.2), Theorem 5.1 implies the following statement.

Theorem 5.2. *The functions (5.17) form an orthonormal complete basis in $\mathcal{L}^2(\mathbb{Q}_p^n)$ (p -adic wavelet basis).*

Corollary 5.2. *The functions*

$$\tilde{\Theta}_{\gamma sa}^{(m)} = F[\Theta_{\gamma sa}^{(m)}](\xi) = p^{n\gamma/2} \chi_p(p^{-\gamma} a \cdot \xi) \Omega(|s + p^{-\gamma} \xi|_p), \quad \xi \in \mathbb{Q}_p^n,$$

form an orthonormal complete basis in $\mathcal{L}^2(\mathbb{Q}_p^n)$, $a = (a_1, \dots, a_n) \in I_p^n$; $s = (s_1, \dots, s_n) \in J_{p;m}^n$; $m = (m_1, \dots, m_n)$, $m_j \geq 1$ is a fixed positive integer, $j = 1, 2, \dots, n$.

The proof follows from Theorem 5.2, formula (6.3) (see below) and the Parseval formula [30, VII,(4.1)].

6. p -ADIC WAVELETS AS EIGENFUNCTIONS OF PSEUDO-DIFFERENTIAL OPERATORS

6.1. Pseudo-differential operators. As mentioned above, the one-dimensional Kozyrev wavelets (5.5) introduced in [20] is a particular case of the wavelets (5.3) for $m = 1$. Moreover, in [20] S. V. Kozyrev proved that his wavelets (5.5) are eigenfunctions of the one-dimensional Vladimirov operator D^α for $\alpha > 0$:

$$D^\alpha \theta_{\gamma ja}(x) = p^{\alpha(1-\gamma)} \theta_{\gamma ja}(x), \quad x \in \mathbb{Q}_p,$$

where $\gamma \in \mathbb{Z}$, $a \in I_p$, $j = 1, 2, \dots, p-1$. Later, it was proved in [3, 4.4.] that in fact, the Kozyrev wavelets (5.5) are eigenfunctions of the Vladimirov operator for any α , i.e., the above formula holds for all $\alpha \in \mathbb{C}$.

Now we prove that n -dimensional wavelets (5.17) are eigenfunctions for a class of pseudo-differential operators (4.1), which includes the Taibleson fractional operator (4.10), (4.6).

Theorem 6.1. *Let A be a pseudo-differential operator with a symbol $\mathcal{A}(\xi) \in \mathcal{E}(\mathbb{Q}_p^n \setminus \{0\})$. Then the n -dimensional p -adic wavelet (5.17) is an eigenfunction of A if and only if*

$$(6.1) \quad \mathcal{A}(p^\gamma(-s + \eta)) = \mathcal{A}(-p^\gamma s), \quad \forall \eta \in \mathbb{Z}_p^n,$$

where $\gamma \in \mathbb{Z}$, $j \in J_{p;m}^n$, $a \in I_p^n$. Here the eigenvalue $\lambda = \mathcal{A}(-p^\gamma s)$, i.e.,

$$A\Theta_{\gamma sa}^{(m)}(x) = \mathcal{A}(-p^\gamma s)\Theta_{\gamma sa}^{(m)}(x).$$

Proof. Let $\Theta_s^{(m)}(x) = \chi_p(s \cdot x) \Omega(|x|_p)$; $x \in \mathbb{Q}_p^n$; $s = (s_1, \dots, s_n) \in J_{p;m}^n$, $s_k \in J_{p;m_k}^n$, $k = 1, 2, \dots, n$. Using (2.2), (2.10), (2.9), we have

$$\begin{aligned} F[\Theta_s^{(m)}(x)](\xi) &= F\left[\prod_{k=1}^n \chi_p(x_k s_k) \Omega(|x_k|_p)\right](\xi) = \prod_{k=1}^n F\left[\Omega(|x_k|_p)\right](\xi_k + s_k|_p) \\ (6.2) \quad &= \prod_{k=1}^n \Omega(|\xi_k + s_k|_p) = \Omega(|\xi + s|_p), \quad \xi \in \mathbb{Q}_p^n. \end{aligned}$$

Here, in view of (2.2), $\Omega(|\xi + s|_p) = \Omega(|\xi_1 + s_1|_p) \times \dots \times \Omega(|\xi_n + s_n|_p)$.

According to (5.2), $|s_k|_p = p^{m_k}$, i.e., $\Omega(|\xi_k + s_k|_p) \neq 0$ only if $\xi_k = -s_k + \eta_k$, where $\eta_k \in \mathbb{Z}_p$, $s_k \in J_{p;m_k}$, $k = 1, 2, \dots, n$. Thus $\xi = -s + \eta$, where $\eta \in \mathbb{Z}_p^n$, $s \in J_{p;m}^n$, and in view of (2.1), $|\xi|_p = p^{\max\{m_1, \dots, m_n\}}$.

In view of formulas (5.17), (6.2), (2.9), we have

$$\begin{aligned} F[\Theta_{\gamma sa}^{(m)}(x)](\xi) &= p^{-n\gamma/2} F[\Theta_s^{(m)}(p^\gamma x - a)](\xi) \\ (6.3) \quad &= p^{n\gamma/2} \chi_p(p^{-\gamma} a \cdot \xi) \Omega(|s + p^{-\gamma} \xi|_p). \end{aligned}$$

Let condition (6.1) be satisfied. Then (4.1), (6.3) imply

$$\begin{aligned} A\Theta_{\gamma sa}^{(m)}(x) &= F^{-1}[\mathcal{A}(\xi) F[\Theta_{\gamma sa}^{(m)}](\xi)](x) \\ (6.4) \quad &= p^{n\gamma/2} F^{-1}[\mathcal{A}(\xi) \chi_p(p^{-\gamma} a \cdot \xi) \Omega(|s + p^{-\gamma} \xi|_p)](x). \end{aligned}$$

Making the change of variables $\xi = p^\gamma(\eta - s)$ and using (2.10), we obtain

$$\begin{aligned} A\Theta_{\gamma sa}^{(m)}(x) &= p^{-n\gamma/2} \int_{\mathbb{Q}_p^n} \chi_p(-(p^\gamma x - a) \cdot (\eta - s)) \mathcal{A}(p^\gamma(\eta - s)) \Omega(|\eta|_p) d^n \eta \\ &= p^{-n\gamma/2} \mathcal{A}(-p^\gamma s) \chi_p(s \cdot (p^\gamma x - a)) \int_{B_0^n} \chi_p(-(p^\gamma x - a) \cdot \eta) d^n \eta \\ &= \mathcal{A}(-p^\gamma s) \Theta_{\gamma sa}^{(m)}(x). \end{aligned}$$

Consequently, $A\Theta_{\gamma sa}^{(m)}(x) = \lambda \Theta_{\gamma sa}^{(m)}(x)$, where $\lambda = \mathcal{A}(-p^\gamma s)$.

Conversely, if $A\Theta_{\gamma sa}^{(m)}(x) = \lambda \Theta_{\gamma sa}^{(m)}(x)$, $\lambda \in \mathbb{C}$, then, using (4.1), (6.3), (6.4), we have

$$(\mathcal{A}(\xi) - \lambda) \Omega(|s + p^{-\gamma} \xi|_p) = 0, \quad \xi \in \mathbb{Q}_p^n.$$

The latter equation has a nontrivial solution only if $s + p^{-\gamma} \xi = \eta$, $\eta \in \mathbb{Z}_p^n$, i.e., $\xi = p^\gamma(-s + \eta)$ and $\lambda = \mathcal{A}(p^\gamma(-s + \eta))$ for any $\eta \in \mathbb{Z}_p^n$. Thus $\lambda = \mathcal{A}(-p^\gamma s)$, and, consequently, (6.1) holds.

The proof of the theorem is complete. \square

The following particular statement was proved in [3].

Corollary 6.1. ([3]) *Let A be a homogeneous pseudo-differential operator with a symbol $\mathcal{A}(\xi) \in \mathcal{E}(\mathbb{Q}_p^n \setminus \{0\})$ of degree π_β . Then the n -direct product $\Theta_{\gamma ja}(x)$ of the one-dimensional Kozyrev p -adic wavelets (5.5), i.e., the n -dimensional p -adic wavelet (5.17) $\Theta_{\gamma sa}^{(1)}$ (for $m = 1$) is an eigenfunction of A if and only if*

$$(6.5) \quad \mathcal{A}(-p^{-1}j + \eta) = \mathcal{A}(-p^{-1}j), \quad \forall \eta \in \mathbb{Z}_p^n,$$

where $\gamma \in \mathbb{Z}$; $a \in I_p^n$; $j = (j_1, \dots, j_n)$, $j_k = 1, 2, \dots, p-1$, $k = 1, 2, \dots, n$. Here the eigenvalue $\lambda = p^{(1-\beta)\gamma} \mathcal{A}(-p^{-1}j)$, i.e.,

$$A\Theta_{\gamma ja}(x) = p^{(1-\beta)\gamma} \mathcal{A}(-p^{-1}j) \Theta_{\gamma ja}(x).$$

6.2. The Taibleson fractional operator. As mentioned above, the Taibleson fractional operator D_x^β is homogeneous of degree β (see Definition 2.1) and has a symbol $\mathcal{A}(\xi) = |\xi|_p^\beta$, which satisfies the condition (6.1)

$$\mathcal{A}(p^\gamma(-s+\eta)) = |p^\gamma(-s+\eta)|_p^\beta = p^{-\beta\gamma} |-s|_p^\beta = p^{\beta(\max\{m_1, \dots, m_n\} - \gamma)} = \mathcal{A}(-p^\gamma s)$$

for all $\eta \in \mathbb{Z}_p^n$. Thus according to Theorem 6.1, the n -dimensional p -adic wavelet (5.17) is an eigenfunction of D_x^β :

$$(6.6) \quad D_x^\beta \Theta_{\gamma sa}^{(m)}(x) = p^{\beta(\max\{m_1, \dots, m_n\} - \gamma)} \Theta_{\gamma sa}^{(m)}(x), \quad \beta \in \mathbb{C}, \quad x \in \mathbb{Q}_p^n,$$

$$\gamma \in \mathbb{Z}, \quad a \in I_p^n, \quad s \in J_{p;m}^n.$$

ACKNOWLEDGMENTS

The authors are greatly indebted to S. V. Kozyrev, V. I. Polischook, and M. A. Skopina for fruitful discussions.

REFERENCES

- [1] S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, Associated homogeneous p -adic distributions, *J. Math. An. Appl.* **313** (2006) 64–83.
- [2] S. Albeverio, A.Yu. Khrennikov, V. M. Shelkovich, Associated homogeneous p -adic generalized functions, *Dokl. Ross. Akad. Nauk* **393** no. 3 (2003), 300–303. English transl. in *Russian Doklady Mathematics*. **68** no. 3 (2003) 354–357.
- [3] S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, Harmonic analysis in the p -adic Lizorkin spaces: fractional operators, pseudo-differential equations, p -adic wavelets, Tauberian theorems, *Journal of Fourier Analysis and Applications*, Vol. 12, Issue 4, (2006), 393–425.
- [4] S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, Pseudo-differential operators in the p -adic Lizorkin space, *p -Adic Mathematical Physics. 2-nd International Conference*, Belgrade, Serbia and Montenegro, 15 – 21 September 2005, Eds: Branko Dragovich, Zoran Rakic, Melville, New York, 2006, AIP Conference Proceedings – March 29, 2006, Vol. 826, Issue 1, pp. 195–205.
- [5] S. Albeverio, A.Yu. Khrennikov, V.M. Shelkovich, p -Adic wavelets and linear and semi-linear evolutionary pseudo-differential equations. (In preparation)
- [6] I.Ya. Aref'eva, B.G. Dragovic, and I.V. Volovich On the adelic string amplitudes, *Phys. Lett.* **B 209** no. 4 (1998) 445–450.
- [7] V.A. Avetisov, A.H. Bikulov, S.V. Kozyrev, and V.A. Osipov, p -Adic models of ultrametric diffusion constrained by hierarchical energy landscapes, *J. Phys. A: Math. Gen.* **12** (2002) 177–189.
- [8] J.J. Benedetto, and R.L. Benedetto, A wavelet theory for local fields and related groups, *The Journal of Geometric Analysis* **3** (2004) 423–456.
- [9] R.L. Benedetto, Examples of wavelets for local fields, *Wavelets, Frames, and operator Theory*, (College Park, MD, 2003), *Am. Math. Soc.*, Providence, RI, (2004), 27–47.
- [10] A.H. Bikulov, and I.V. Volovich, p -Adic Brownian motion, *Izvestia Akademii Nauk, Seria Math.* **61** no. 3 (1997) 537–552.
- [11] I.M. Gel'fand and G.E. Shilov, Generalized functions. vol 1: Properties and operations. New York, Acad. Press, 1964.
- [12] I.M. Gel'fand, M.I. Graev and I.I. Piatetskii-Shapiro, Generalized functions. vol 6: Representation theory and automorphic functions. Nauka, Moscow, 1966.

- [13] A. Khrennikov, *p*-Adic valued distributions in mathematical physics. Kluwer Academic Publ., Dordrecht, 1994.
- [14] A. Khrennikov, *Non-archimedean analysis: quantum paradoxes, dynamical systems and biological models*. Kluwer Academic Publ., Dordrecht, 1997.
- [15] A. Khrennikov, *Information dynamics in cognitive, psychological, social and anomalous phenomena*. Kluwer Academic Publ., Dordrecht, 2004.
- [16] A.Yu. Khrennikov, and S.V. Kozyrev, Wavelets on ultrametric spaces, *Applied and Computational Harmonic Analysis* **19** (2005) 61–76.
- [17] A.Yu. Khrennikov, and S.V. Kozyrev, Pseudodifferential operators on ultrametric spaces and ultrametric wavelets, *Izvestia Akademii Nauk, Seria Math.* **69** no. 5 (2005) 133–148.
- [18] A. Yu. Khrennikov, and S. V. Kozyrev, Localization in space for free particle in ultrametric quantum mechanics, *Dokl. Ross. Akad. Nauk* **411** no. 3 (2006) 316–322. English transl. in *Russian Doklady Mathematics*. **74** no. 3 (2006) 906–911.
- [19] A.N. Kochubei, *Pseudo-differential equations and stochastics over non-archimedean fields*, Marcel Dekker. Inc. New York, Basel, 2001.
- [20] S.V. Kozyrev, Wavelet analysis as a *p*-adic spectral analysis, *Izvestia Akademii Nauk, Seria Math.* **66** no. 2 (2002) 149–158.
- [21] S.V. Kozyrev, *p*-Adic pseudodifferential operators: methods and applications, *Proc. Steklov Inst. Math.* **245**, Moscow (2004) 154–165.
- [22] S.V. Kozyrev, *p*-Adic pseudodifferential operators and *p*-adic wavelets, *Theor. Math. Physics* **138**, no. 3 (2004) 1–42.
- [23] S.V. Kozyrev, V.Al. Osipov, V.C. A.Avetisov, Nondegenerate ultrametric diffusion, *J. Math. Phys.* **46** no. 6 (2005) 15 pp.
- [24] P.I. Lizorkin, Generalized Liouville differentiation and the functional spaces $L_p^r(E_n)$. Imbedding theorems, (Russian) *Mat. Sb. (N.S.)* **60**(102) (1963) 325–353.
- [25] P.I. Lizorkin, Operators connected with fractional differentiation, and classes of differentiable functions, (Russian) *Studies in the theory of differentiable functions of several variables and its applications, IV. Trudy Mat. Inst. Steklov.* Vol. 117 (1972), 212–243.
- [26] S.G. Samko, *Hypersingular integrals and their applications*. Taylor & Francis, London, 2002.
- [27] S.G. Samko, A.A. Kilbas, and O.I. Marichev, *Fractional integrals and derivatives and some of their applications*. Minsk, Nauka i Tekhnika, 1987 (in Russian); English translation: *Fractional integrals and derivatives. Theory and applications*, Gordon and Breach, London, 1993.
- [28] M.H. Taibleson, Harmonic analysis on *n*-dimensional vector spaces over local fields. I. Basic results on fractional integration, *Math. Annalen* **176** (1968) 191–207.
- [29] M.H. Taibleson, *Fourier analysis on local fields*. Princeton University Press, Princeton, 1975.
- [30] V.S. Vladimirov, I.V. Volovich and E.I. Zelenov, *p*-Adic analysis and mathematical physics. World Scientific, Singapore, 1994.
- [31] V.S. Vladimirov, I.V. Volovich, *p*-Adic quantum mechanics, *Commun. Math. Phys.* **123** (1989) 659–676.
- [32] I.V. Volovich, *p*-Adic string, *Class. Quant. Grav.* **4** (1987) L83–L87.
- [33] W.A. Zuniga-Galindo, Pseudo-differential equations connected with *p*-adic forms and local zeta functions, *Bull. Austral. Math. Soc.* **70** no. 1 (2004) 73–86.
- [34] W.A. Zuniga-Galindo, Fundamental solutions of pseudo-differential operators over *p*-adic fields, *Rend. Sem. Mat. Univ. Padova* **109** (2003) 241–245.

INTERNATIONAL CENTER FOR MATHEMATICAL MODELLING IN PHYSICS AND COGNITIVE SCIENCES MSI, VÄXJÖ UNIVERSITY, SE-351 95, VÄXJÖ, SWEDEN. PHONE: +046 (470) 708790 FAX: +046 (470) 84004

E-mail address: `andrei.khrennikov@msi.vxu.se`

DEPARTMENT OF MATHEMATICS, ST.-PETERSBURG STATE ARCHITECTURE AND CIVIL ENGINEERING UNIVERSITY, 2 KRASNOARMEISKAYA 4, 190005, ST. PETERSBURG, RUSSIA. PHONE: +7 (812) 2517549 FAX: +7 (812) 3165872

E-mail address: `shelkv@vs1567.spb.edu`